

# The quark star in the Quark Mean Field approach.

Ryszard Mańka<sup>\*</sup> and Grzegorz Przybyła<sup>†</sup>

*University of Silesia, Institute of Physics,  
Katowice 40007, ul. Uniwersytecka 4, Poland*

(Dated: February 8, 2008)

## Abstract

The quark star in the quark mean field (QMF) extended to include the strange mesons interaction is examined. The quark star properties in this model are studied.

arXiv:nucl-th/0007071v1 30 Jul 2000

---

<sup>\*</sup>Electronic address: [manka@us.edu.pl](mailto:manka@us.edu.pl); URL: [www.cto.us.edu.pl/~manka](http://www.cto.us.edu.pl/~manka); Presented at the workshop *Physics of Neutron Star Interiors* in the European Center for Theoretical Studies in Nuclear Physics and related Areas ECT\* in Trento, Italy, 2000.

<sup>†</sup>Electronic address: [przybyla@phys.server.us.edu.pl](mailto:przybyla@phys.server.us.edu.pl)

## Introduction

Recent theoretical studies show that the properties of nuclear matter can be described nicely in terms of the Relativistic Mean Field Theory (RMF) [1, 2, 3]. Guichon proposed an interesting model on the change of the nucleon properties in nuclear matter (QMF or quark-meson coupling model (QMC)) [4]. The model construction mimics the relativistic mean field theory, where the scalar  $\sigma$  and the vector meson  $\omega$  fields couple not with nucleons but directly with quarks. The quark mass has to change from its bare mass due to the coupling to the  $\sigma$  meson. In this work we shall investigate the quark matter within the Relativistic Mean Field theory motivated by the Nambu-Jona-Lasinio (NJL) model [5][6][7]. The three-flavor NJL model has been discussed by many authors, e.g. [8]. To include the strange quark the QMF model is extended by incorporating an additional pair of the strange meson fields which couple only to the  $s$  quark [9]. It is very interesting to construct the Guichon model, where the nucleon is described in terms of constituent quarks, which couple with mesons and gluons. This model (we refer it as the quark mean field (QMF) model was named in [10]). The meson fields act on quarks inside a nucleon or the neutron or quark star and change the bulk nuclear properties. The QMF model predicts an increasing of the nuclear size and a reduction of the nucleon mass in the nuclear environment of the star.

The properties of strange quark matter (SQM) and the conjecture that SQM could be the absolute ground state of strongly interacting matter [11, 12] have attracted much attention in nuclear physics and astrophysics. In 1984 Farhi and Jaffe [13] found that SQM is stable compared with an  $^{56}\text{Fe}$ -nucleus. Still the bag model calculations play a key role in this field [14].

In the framework of the NJL model Babulla and Oertel [6] have showed that due to a large constituent quark the strange star is not absolutely stable. The quark-meson coupling will reduce the effective quark mass what may make the existence of the strange star more possible.

The aim of this work is to examine the influence of the quark-meson coupling model on the quark star or neutron star having the quarks core.

## The Quark Mean Field Theory

The fields of the model QMF for  $\sigma$ ,  $\omega$  and  $\rho$ -mesons are denoted as  $\varphi$ ,  $\omega_\mu$ ,  $\rho_\mu$ . To reproduce the observed strongly attractive  $\Lambda\Lambda$  interaction two additional meson fields are added, the scalar  $f_0$  and meson fields  $\phi$  denoted as the fields  $\varphi_*$  and  $\phi_\mu$ . The Lagrange density function for this model has the following form

$$\begin{aligned}\mathcal{L} = & -\frac{1}{4}R_{\mu\nu}^a R^{a\mu\nu} - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{1}{2}M_\omega^2 \omega_\mu \omega^\mu + \frac{1}{2}M_\rho^2 \rho_\mu^a \rho^{a\mu} \\ & \frac{1}{2}\partial_\mu \varphi \partial^\mu \varphi - U(\varphi) + \frac{1}{4}c_3(\omega_\mu \omega^\mu)^2 + \\ & i \sum_{f=1}^2 \bar{L}_f \gamma^\mu \partial_\mu L_f - \sum_{f=1}^2 g_f (\bar{L}_f H e_{Rf} + h.c.) + \\ & i \bar{q} \gamma^\mu D_\mu q - \bar{q} (m_Q - g_s^q \varphi I_2 - g_{s*}^q \varphi_* I_3) q + \\ & \frac{1}{2} \partial_\mu \varphi_* \partial^\mu \varphi_* - \frac{1}{2} m_{s*}^2 \varphi_*^2 - \frac{1}{4} \phi_{\mu\nu} \phi^{\mu\nu} + \frac{1}{2} M_\phi^2 \phi_\mu \phi^\mu.\end{aligned}\tag{1}$$

The field stress tensor of the vector mesons have the following structure:

$$R_{\mu\nu}^a = \partial_\mu \rho_\nu^a - \partial_\nu \rho_\mu^a + g \varepsilon_{abc} \rho_\mu^b \rho_\nu^c \tag{2}$$

$$F_{\mu\nu} = \partial_\mu \omega_\nu - \partial_\nu \omega_\mu \tag{3}$$

$$G_{\mu\nu}^a = \partial_\mu W_\nu^a - \partial_\nu W_\mu^a + g f_{abc} W_\mu^b W_\nu^c$$

$$\phi_{\mu\nu} = \partial_\mu \phi_\nu - \partial_\nu \phi_\mu \tag{4}$$

The covariant derivative acts on baryons as follows:

$$D_\mu = \partial_\mu + \frac{1}{2} i g_\rho^q \rho_\mu^a \tau^a + i g_\omega^q \omega_\mu I_2 + i g_\phi^q \phi_\mu I_3 \tag{5}$$

The potential in RMF [1, 15, 16] is given by

$$U(\varphi) = B_0 + \frac{1}{2} m_s^2 \varphi^2 + \frac{1}{3} g_2 \varphi^3 + \frac{1}{4} g_3 \varphi^4 \tag{6}$$

The fermion fields are composed of quarks and electrons, muons and neutrinos

$$q = \begin{pmatrix} u \\ d \\ s \end{pmatrix}, \quad I_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad I_3 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \tau^a = \begin{pmatrix} \sigma^a & 0 \\ 0 & 0 \end{pmatrix} \tag{7}$$

| Parameter                                                    | TM1[15, 16]              |
|--------------------------------------------------------------|--------------------------|
| $M$                                                          | 938 MeV                  |
| $M_\omega$                                                   | 783 MeV                  |
| $M_\rho$                                                     | 770 MeV                  |
| $m_s$                                                        | 511.198 MeV              |
| $m_{s*}$                                                     | 975 MeV                  |
| $M_\phi$                                                     | 1020 MeV                 |
| $m_Q = m_u^* = m_d^* = m_s^*$                                | 310.6 MeV [6]            |
| $m_Q = m_u^* = m_d^* \neq m_s^*$                             | 367.7 MeV, 549.5 MeV [7] |
| $g_3$                                                        | 7.2325 fm <sup>-1</sup>  |
| $g_4$                                                        | 0.6183                   |
| $g_s = g_{\sigma N} = 3g_s^q$                                | 10.0289                  |
| $g_\omega = g_{\omega N} = 3g_\omega^q$                      | 12.6139                  |
| $g_\rho = g_{\rho N} = g_\rho^q$                             | 4.6322                   |
| $g_{s*} = g_{\sigma*\Lambda} = 2/3 g_s$                      | $g_{s*} = 3g_{s*}^q$     |
| $g_\phi = g_{\phi\Lambda} = \sqrt{2}/3 g_\omega = 3g_\phi^q$ | $c_3 = 71.3075$          |

TABLE I: Parameter sets for the Lagrangian function (1).

$$L_1 = \begin{bmatrix} \nu_e \\ e^- \end{bmatrix}_L, \quad L_2 = \begin{bmatrix} \nu_\mu \\ \mu^- \end{bmatrix}_L, \quad e_{Rf} = (e_R^-, \mu_R^-). \quad (8)$$

$m_s$ ,  $M_\omega$ ,  $M_\rho$  are masses assigned to the mesons fields Here  $q$  denotes a quark field with three flavors,  $u$ ,  $d$  and  $s$ , and three colors. The Lagrangian function includes also the nonlinear term  $\frac{1}{4}c_3(\omega_\mu\omega^\mu)^2$ . We start with equal dynamical quark masses  $m_Q = m_u^* = m_d^* = m_s^* = 310.6 \text{ MeV}$  (the  $SU(3)$ -symmetric quark matter with starting  $m_{0,u} = m_{0,d} = m_{0,s} = 0$ )[6] and a bag pressure  $B_0 = 57.3 \text{ MeV/fm}^3$  originated from the Nambu-Johna-Lasinio model

$$L_{NJL} = i\bar{q}\gamma^\mu D_\mu q - \bar{q}m_0q + G \sum_{a=0}^8 \{(\bar{q}\lambda^a q)^2 + (i\bar{q}\lambda^a \gamma^5 q)^2\}. \quad (9)$$

We restrict ourselves to the isospin  $SU(2)$  symmetric case,  $m_0^u = m_0^d$ , so

$$m_0 = \begin{pmatrix} m_{0,u} & & \\ & m_{0,d} & \\ & & m_{0,s} \end{pmatrix}$$

In the quark massless limit the system has a  $U(3)_L \times U(3)_R$  chiral symmetry. The model is not renormalizable and we have to specify a regularization scheme for divergent integrals. For simplicity we use a sharp cut-off  $\Lambda$  in 3-momentum space. Besides  $\Lambda$  we have to fix the coupling constants  $G$  and the current quark masses  $m_{0,u}$  and  $m_{0,s}$ . In most of our calculations we will adopt the parameters of [6]  $\Lambda = 602.3$  MeV,  $G\Lambda^2 = 1.835$ . With  $\Lambda$ ,  $G$  and  $K$  as specified above, chiral symmetry is spontaneously broken. The possible existence of deconfined quark matter in the interior of neutron stars using the NJL model was examined in paper [7]. In the work [7] the non-symmetrical quark masses  $m_Q = \{m_u^* = m_d^* = 367.7 \text{ MeV}, m_s^* = 549.5 \text{ MeV}\}$  are obtained (with starting  $m_{0,u} = m_{0,d} = 5.5$  MeV,  $m_{0,s} = 140.7$  MeV).

The Euler equation for  $\Phi_A = \{\varphi, \varphi_*, \omega_\mu, \rho_\mu, \phi_\mu, q\}$  fields are

$$\square\varphi = m_s^2\varphi + g_2\varphi^2 + g_3\varphi^3 - g_s^q\bar{q}I_2q \quad (10)$$

$$\square\varphi_* = m_{s*}^2\varphi_* - g_s^q\bar{q}I_3q \quad (11)$$

$$-\partial_\mu F^{\mu\nu} = M_\omega^2\omega^\nu + c_3(\omega_\mu\omega^\mu)\omega^\nu - g_\omega^q J_{SU(2)}^\nu \quad (12)$$

$$-\partial_\mu \phi^{\mu\nu} = M_\phi^2\phi^\nu - g_\phi^q J_{U(1)}^\nu \quad (13)$$

where

$$J_{SU(2)}^\nu = \bar{q}\gamma^\nu I_2q \quad (14)$$

is the  $SU(2)$  up and down quark current while

$$J_{U(1)}^\nu = \bar{q}\gamma^\nu I_3q \quad (15)$$

is the  $U(1)$  strange quark current. The nonabelian gauge field  $\rho^{\nu,a}$  obeys

$$-D_\mu R^{\mu\nu,a} = M_\rho^2 \rho^{\nu,a} - g_\rho^q J_3^\nu \quad (16)$$

where

$$J_3^\nu = \frac{1}{2}\bar{q}\gamma^\nu \tau^3 q \quad (17)$$

is the isospin SU(2) current. In the system we have conservation of “nucleon” charge

$$Q_N = \int d^3x J_{SU(2)}^0$$

the isospin charge

$$Q_3 = \int d^3x J_3^0$$

and the “hyperon” charge

$$Q_H = \int d^3x J_{U(1)}^0$$

The last is the Dirac equation

$$i\gamma^\mu D_\mu q - (m_Q - g_s^q \varphi I_2 - g_{s*}^q \varphi_* I_3)q = 0. \quad (18)$$

The physical system is totally defined by the thermodynamic potential [17]

$$\Omega = -kT \ln \text{Tr}(e^{-\beta(H - \mu_N Q_N - \mu_3 Q_3 - \mu_H Q_H)}) \quad (19)$$

where H is the Hamiltonian of the physical system

$$H = \sum_A \int d^3x \{ \partial_0 \Phi_A \pi_\Phi^A - \mathcal{L} \} \quad (20)$$

and  $\pi^A = \frac{\partial \mathcal{L}}{\partial(\partial_0 \Phi_A)}$  is a momentum connected to the field  $\Phi_A$ . The fields  $\Phi_A = \{\varphi, \varphi_*, \omega_\mu, \rho_\mu, \phi_\mu, q\}$  denote all fields in the system. In this paper we shall use the effective potential approach build using the Bogolubov inequality [15, 18]

$$\Omega \leq \Omega_1 = \Omega_0(m_B, m_F) + \langle H - H_0 \rangle_0 \quad (21)$$

$\Omega_0$  is the thermodynamic potential of the trial system as effectively free quasiparticle system described by the Lagrange function

$$\begin{aligned} \mathcal{L}_0 = & \frac{1}{2} \partial_\mu \bar{\varphi} \partial^\mu \bar{\varphi} - \frac{1}{2} m_s^2 \bar{\varphi}^2 + \frac{1}{2} \partial_\mu \bar{\varphi}_* \partial^\mu \bar{\varphi}_* - \frac{1}{2} m_s^2 \bar{\varphi}_*^2 + \\ & - \frac{1}{4} \bar{R}_{\mu\nu}^a \bar{R}^{a\mu\nu} - \frac{1}{4} \bar{F}_{\mu\nu} \bar{F}^{\mu\nu} - \frac{1}{4} \bar{\phi}_{\mu\nu} \bar{\phi}^{\mu\nu} \\ & + \frac{1}{2} M_\omega^2 \bar{\omega}_\mu \bar{\omega}^\mu + \frac{1}{2} M_\rho^2 \bar{\rho}_\mu^a \bar{\rho}^{a\mu} + \frac{1}{2} M_\phi^2 \bar{\phi}_\mu \bar{\phi}^\mu + \sum_{f=u,d,s} \bar{q}_f (i\gamma^\mu \bar{D}_\mu - m_{F,f}) q_f \end{aligned} \quad (22)$$

Similar to the general case

$$\bar{R}_{\mu\nu}^a = \partial_\mu \bar{\rho}_\nu^a - \partial_\nu \bar{\rho}_\mu^a$$

and

$$\bar{F}_{\omega,\mu\nu} = \partial_\mu \bar{\omega}_\nu - \partial_\nu \bar{\omega}_\mu.$$

and for gluons

$$\overline{G}_{\mu\nu}^a = \partial_\mu \overline{W}_\nu^a - \partial_\nu \overline{W}_\mu^a$$

We decompose the  $\Phi_A$  field into two components, the effectively free quasiparticle field  $\tilde{\Phi}_A$  and the classical boson condensate  $\xi_A$

$$\Phi_A = \tilde{\Phi}_A + \xi_A \quad (23)$$

In the case of the RMF model we have

$$\varphi = \overline{\varphi} + \sigma \quad (24)$$

$$\varphi_* = \overline{\varphi}_* + \sigma_* \quad (25)$$

$$\omega_\mu = \overline{\omega}_\mu + w_\mu, \quad w_\mu = \delta_{\mu,0} w \quad (26)$$

$$\rho_\mu^a = \overline{\rho}^a + r_\mu^a, \quad r_\mu^a = \delta^{a,3} \delta_{\mu,0} r \quad (27)$$

$$\phi_\mu = \overline{\phi}_\mu + w_{\mu*}, \quad w_{\mu*} = \delta_{\mu,0} w_* \quad (28)$$

The  $\xi_A = \{\sigma, w, r, w_*\}$  field will be treated as the variational parameters in the effective potential. Also the boson and fermion mass  $m_B, m_F$  will be treated as the variational parameters. The covariant derivative for the trial system is

$$\overline{D}_\mu = \partial_\mu + \frac{1}{2} i g_\rho^q r_\mu^a \sigma^a + i g_\omega^q w_\mu + i g_\phi^q w_{\mu*} \quad (29)$$

This introduce the homogenous fermion interaction with boson condensate  $w_\mu, r_\mu^a, w_{\mu*}$ . The fermion quasiparticle will obey the Dirac equation

$$(i\gamma^\mu \overline{D}_\mu - m_F)\psi_f = 0 \quad (30)$$

The constant condensate  $w, r$  simply shift the chemical potential from  $\mu_i = \mu_i^0$  (when  $w = r = 0$ ) to

$$\mu_u = \mu_u^0 + \frac{1}{2} g_\rho^q r - g_\omega^q w \quad (31)$$

$$\mu_d = \mu_d^0 - \frac{1}{2} g_\rho^q r - g_\omega^q w \quad (32)$$

$$\mu_s = \mu_s^0 - g_\phi^q w_* \quad (33)$$

where  $\mu_u = \mu + \frac{1}{2}\mu_3$  and  $\mu_d = \mu - \frac{1}{2}\mu_3$ .

Quarks and electrons are in  $\beta$ -equilibrium which can be described as a relation among their chemical potentials

$$\mu_d = \mu_u + \mu_e = \mu_s$$

where  $\mu_u$ ,  $\mu_d$ ,  $\mu_s$  and  $\mu_e$  stand for quarks and electron chemical potentials respectively. If the electron Fermi energy is high enough (greater than the muon mass) in the neutron star matter muons start to appear as a result of the following reaction

$$d \rightarrow u + e^- + \bar{\nu}_e$$

$$s \rightarrow u + \mu^- + \bar{\nu}_\mu$$

In a pure quark state the star should to be charge neutral. This gives us an additional constraint on the chemical potentials

$$\frac{2}{3}n_u - \frac{1}{3}n_d - \frac{1}{3}n_s - n_e = 0. \quad (34)$$

where  $n_f$  ( $f \in u, d, s$ ),  $n_e$  the particle densities of the quarks and the electrons, respectively. The EOS can now be parameterized by only one chemical potential, say  $\mu_u = Mx$ . Variation

$$\frac{\partial f_1}{\partial m_F} = 0$$

with respect to the trial system  $L_0$  gives

$$m_{F,u,d} = m_Q - g_s^q \sigma = M\delta_{u,d} \quad (35)$$

$$m_{F,s} = m_Q - g_{s*}^q \sigma = M\delta_s \quad (36)$$

In the local equilibrium inside the star the free energy reaches the minimum at  $\sigma$ .

The same result may be achieved calculating the averages of the equation of motions (10) for the effective system  $\mathcal{L}_0$ . In the mean field approximation the meson field operators are replaced by their expectation values. We also consider the isotropic system at rest. As  $\langle \bar{q}q \rangle_0$  (calculated with respect to  $\mathcal{L}_0$  system) depends on the effective quark mass  $m_F$  (or  $\sigma$ ,  $\sigma_*$ ) the equation (10) is highly nonlinear also with respect to  $\sigma$ .

Calculation base on the relation

$$\frac{\partial f_F}{\partial m_{F,f}} = \langle \bar{q}q \rangle_0$$






FIG. 1: The effective quark masses.

gives

$$\langle \bar{q}_f q_f \rangle_0 = \frac{m_F}{\pi^2} \int_0^\infty \frac{p^2 dp}{\sqrt{p^2 + m_F^2}} \left\{ \frac{1}{\exp(\beta(\epsilon_p - \mu_f)) + 1} + \frac{1}{\exp(\beta(\epsilon_p + \mu_f)) + 1} \right\}. \quad (37)$$

The quantum average  $\langle \bar{q}_f q_f \rangle_0$  depends on the quark chemical potentials (31,32). In the result the effective quark effective mass  $m_F$  also will be dependent on the chemical potentials. In the result the solution  $\sigma$  of the equation (10) also will be dependent. The same situation will consider other fields.

The effective quarks mass  $m_{F,f}$  (or  $\delta_f = m_{F,f}/M$ ) dependence on the dimensionless Fermi momentum  $x_F$  is presented on the Fig.1. To calculate the properties of the quark star we need the energy-momentum tensor. We define the density of energy and pressure by the energy - momentum tensor

$$\langle T_{\mu\nu} \rangle = (P + \epsilon) u_\mu u_\nu - P g_{\mu\nu}$$

where  $u_\mu$  is a unite vector ( $u_\mu u^\mu = 1$ ). Similar to paper [15, 19] we have introduced the dimensionless “Fermi” momentum even at finite temperature which exactly corresponds to the Fermi momentum at zero temperature. Both  $\epsilon_F$  and  $P_F$  depend on the quark chemical




FIG. 2: The quark and neutron (TM1) and hyperon equation of state.

potential  $\mu$  or Fermi momentum  $x_F$ . This parametric dependence on  $\mu$  (or  $x_F$ ) defines the equation of state. The various equations of state for different parameters sets is presented on Fig. 2.

### The quark star properties

The numerical results describing the structure of neutron or quark star are based on the relativistic mean field theory are now presented. It is possible to describe a static spherical star solving the OTV equation.

$$\frac{dP(r)}{dr} = -\frac{G}{r^2}(\rho(r) + \frac{P(r)}{c^2}) \frac{(m(r) + \frac{4\pi}{c^2}P(r)r^3)}{(1 - \frac{2Gm(r)}{c^2r})} \quad (38)$$

$$\frac{dm(r)}{dr} = 4\pi r^2 \rho(r) \quad (39)$$

Having solved the OTV equation the pressure  $p(r)$ , mass  $m(r)$  and density  $\rho(r)$  were obtained. To obtain the total radius  $R$  of the star the fulfillment of the condition  $p(R) = 0$  is necessary. This allows to determine the total gravitational mass of the star  $M(R)$ . The  $M(R)$  for the quark star is presented on the Fig. 3. The  $R(\rho)$  dependence for the quark star is presented on the Fig. 4. For the quark star with the central density  $\rho_c = 2 \cdot 10^{15} \text{ g/cm}^3$  the




FIG. 3: The  $M(R)$  dependence for the quark star.




FIG. 4: The  $R(\rho)$  dependence for the quark star.

star profile in the mean field approach is presented on the Figs. 5,6,8. The effective quark chemical potential are presented on the Fig. 5.

The quark and electron Fermi momentum inside the star is presented on Fig. 6. This Fermi momentum distribution comes from the quark dispersion relation presented on the




FIG. 5: The effective quark chemical potential for the quark star with the central density  $\rho_c = 2 \cdot 10^{15} \text{ g/cm}^3$

Fig. 7. The meson field  $\omega_\mu$  and  $\phi_\mu$  profile inside the star is presented on the Fig. 8. The biggest contributions to the energy density come from the quarks, the gauge boson field  $\omega_\mu$ ,  $\phi_\mu$  and scalar boson field  $\varphi$ ,  $\varphi_*$ . The quark effective mass profile inside the star is presented on Fig. 9. This difference between strange and up, down quarks deepens inside the star. Its influences on the quark partial fraction defined as

$$X_f = \frac{n_f}{(n_u + n_d + n_s)}, \quad (40)$$

where  $f = (u, d, s)$ , presented on the Fig. 10. We see that both pure neutron star ( $X_d = 2/3$ ,  $X_u = 1/3$ ) and strange star ( $X_u = X_d = X_s$ ) are rather mathematical limits. The electron distribution inside the quark star is presented on the Fig. 11.

In the flat spacetime the preferred configuration is realized for the system with minimum of free energy. As pressure is defined as

$$P = -\frac{\partial F}{\partial V},$$

this means that the stable configuration should have bigger pressure. According to the equation of state (Fig. 2) as was pointed in the strange star configuration is unstable. As a consequence there seems to be no chance to find SQM with energies per baryon number




FIG. 6: The quark and electron Fermi momentum inside the star




FIG. 7: The quark Fermi momentum distribution.

lower than in ordinary nuclei. This almost rules out the original idea of absolutely stable SQM.

However, when hyperon are included there is crossing point at

$$\rho_{cs} = 8.92 \cdot 10^{14} \text{ g/cm}^3$$




FIG. 8: The meson field  $\omega_\mu$  and  $\phi_\mu$  profile inside the quark star.




FIG. 9: The quark effective mass profile inside the star.

above which the strange quark core (the Buballa, Oertel parameters set [6]) may appear. This will happen when the star central density  $\rho_c$  exceed  $\rho_{cs}$ . However, Fig.4 shows that the hyperon star is unstable in this region. A mixed hyperon star with strange quark core can be stabilized if the the equation of state will be more softer (including the kaon condensation




FIG. 10: The quark partial fraction inside the star.




FIG. 11: The electron distribution inside the quark star.

[20], for example).

Unfortunately, the strange star with Schetler et al. parameters set [7] is unstable due to a large constituent strange quark mass ( $m_s^* = 549.5 \text{ MeV}$ ). In the flat Minkowski spacetime the nucleon binding energy is presented on Fig.12.




FIG. 12: The nuclear binding energy as the relative baryon number  $u = n/n_0$  ( $n_0 = 0.17 \text{ fm}^{-3}$  for the nuclear symmetric matter).

All these considerations concerning the star stability were done in the flat Minkowski spacetime. However, a neutron or quark star is mainly binded by gravity. They binding energy [21] are equal to

$$\Delta E = c^2 M - c^2 M_B = (c^2 M - c^2 M_p) + (c^2 M_p - c^2 M_B), \quad (41)$$

where

$$M_p = \int_0^R 4\pi r^2 dr \left(1 - \frac{2Gm(r)}{c^2 r}\right)^{-\frac{1}{2}} \rho(r) \quad (42)$$

is the proper mass of the compact object and

$$M = \int_0^R 4\pi r^2 dr \rho(r) \quad (43)$$

is the gravitational mass. The gravitational binding energy for the hyperon and quark star is presented on Fig. 13.  $M_B$  is the baryonic mass defined as

$$M_B = \sum_B m_B N_B \quad (44)$$

where  $m_B$  is the baryon mass and the baryon number is equal to

$$N_B = \int_0^R 4\pi r^2 dr \left(1 - \frac{2Gm(r)}{c^2 r}\right)^{-\frac{1}{2}} n_B(r) \quad (45)$$

The star mass deficit

$$\Delta M = (M_p - M)$$






FIG. 13: The star gravitational binding energy as a function of the star baryon number  $N/N_\odot$  ( $N_\odot = 1.187 \cdot 10^{57}$  is the baryon number of the Sun).




FIG. 14: The star mass deficit calculated in the Sun units ( $M_\odot c^2 = 1.787 \cdot 10^{54} \text{ ergs}$ ) as a function of the star central density.

is presented on Fig.14. The quark star is stable with respect to the gravitational binding energy (Fig. 13). The difference of the gravitational binding energy for the hyperon and quark star is order of

$$\sim 8.9 \cdot 10^{52} \text{ ergs} \quad (46)$$

Such a doze of energy may be released during conversion of the neutron star into quark star. For example, the  $M_B = M_\odot$  hyperon star with the mass  $M \sim 0.92 M_\odot$ , radius

$R \sim 13 \text{ km}$  and central density  $\rho_c \sim 4 \text{ g/cm}^3$  will transform into the quark star with the mass  $M \sim 0.85 M_\odot$ , radius  $R \sim 8.5 \text{ km}$  and central density  $\rho_c \sim 8.9 \text{ g/cm}^3$  (Figs. 13,14). This conversion may explain nature of the gamma-ray bursts [22].

## Conclusion

Quark meson coupling model was designed to describe both the bulk and internal structures of nuclear phenomenology described in terms of the Relativistic Mean Field Theory (RMF) [1]. The hyperon-hyperon interaction mediated by the  $\varphi_*$  and  $\phi_\mu$  mesons is expected to be important for hyperon rich matter present in the cores of neutron star or core of the quark star. The quarks interaction is a bit different for the  $u$ ,  $d$  and  $s$  quarks. This interaction breaks the starting SU(3) flavor symmetry. The real quark star still is dominated by the  $u$ ,  $d$  quarks with smaller presence of the  $s$  quarks. Only in the mathematical limit of the core high density we have the strange star ( $X_u = X_d = X_s$ ) case. The appearance of deconfined quark matter in the center of the neutron star demands soft nuclear equation of state. The quark star occurs stable when the gravitational binding takes into consideration.

## Appendix: The hyperon star in the Relativistic Mean Field Theory

To compare the quark star with neutron one we shall use the RMF model of the neutron star including the  $\Lambda$  hyperon [23]. This is a simple generalization of the paper [15]. The model consists following mesons:  $\varphi$ ,  $\omega_\mu$ ,  $\rho_\mu^a$  with masses  $m_s = 510 \text{ MeV}$ ,  $M_\omega = 783 \text{ MeV}$ ,  $M_\rho = 770 \text{ MeV}$ , and  $\varphi_*$ ,  $\phi_\mu$  with masses  $m_{s*} = 975 \text{ MeV}$ ,  $M_\phi = 1020 \text{ MeV}$ . The fermion fields are composed of protons, neutrons and  $\Lambda$  hyperon

$$\psi = \begin{pmatrix} \psi_p \\ \psi_n \end{pmatrix}, \quad \psi_\Lambda$$

with masses  $M = 938 \text{ MeV}$ ,  $M_\Lambda = 1115.6 \text{ MeV}$ .

The Lagrange function is:

$$L = -\frac{1}{2}\partial_\mu\varphi\partial^\mu\varphi - U(\varphi) - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{2}M_\omega^2\omega_\mu\omega^\mu \quad (47)$$

$$-\frac{1}{4}c_3(\omega_\mu\omega^\mu)^2 - \frac{1}{4}R_{\mu\nu}^a R^{a\mu\nu} - \frac{1}{2}M_\rho^2\rho_\mu^a\rho^{a\mu} \quad (48)$$

$$-\frac{1}{2}\partial_\mu\varphi_*\partial^\mu\varphi_* - \frac{1}{2}m_{s*}^2\varphi_*^2 - \frac{1}{4}\Phi_{\mu\nu}\Phi^{\mu\nu} - \frac{1}{2}M_\phi^2\phi_\mu\phi^\mu \quad (49)$$

$$+i\bar{\psi}\gamma^\mu D_\mu\psi - \bar{\psi}(M - g_{\sigma N}\varphi)\psi \quad (50)$$

$$+i\bar{\psi}_\Lambda\gamma^\mu D_\mu\psi_\Lambda - \bar{\psi}_\Lambda(M_\Lambda - g_{\sigma\Lambda}\varphi - g_{\sigma*\Lambda}\varphi_*)\psi_\Lambda \quad (51)$$

where  $F_{\mu\nu}$  is the stress tensor of the form

$$F_{\mu\nu} = \partial_\mu\omega_\nu - \partial_\nu\omega_\mu,$$

and the tensor  $R_{\mu\nu}^a$  has the form

$$R_{\mu\nu}^a = \partial_\mu\rho_\nu^a - \partial_\nu\rho_\mu^a + g_\rho\varepsilon_{abc}\rho_\mu^b\rho_\nu^c$$

and the scalar potential is

$$U(\varphi) = \frac{1}{2}M_\sigma^2\varphi^2 + \frac{1}{3}g_2\varphi^3 + \frac{1}{4}g_3\varphi^4,$$

where  $c_3$ ,  $g_2$ ,  $g_3$  are constant of the TM1 model [15, 18]. For the hyperon sector

$$\Phi_{\mu\nu} = \partial_\mu\phi_\nu - \partial_\nu\phi_\mu,$$

The covariant derivative are defined now as

$$D_\mu\psi = (\partial_\mu + \frac{1}{2}ig_{\rho N}\rho_\mu^a\sigma^a + ig_{\omega N}\omega_\mu)\psi, \quad (52)$$

$$D_\mu\psi_\Lambda = (\partial_\mu + ig_{\phi\Lambda}\phi_\mu)\psi_\Lambda \quad (53)$$

The coupling constants are

$$g_{\sigma\Lambda} = \frac{2}{3}g_{\sigma N}$$

$$g_{\sigma*\Lambda} = \frac{\sqrt{2}}{3}g_{\sigma N}$$

for SU(6) symmetry, the coupling constants  $g_{\sigma N}$ ,  $g_{\omega N}$  are the same as for the TM1 parameters set [15] and

$$g_{\phi\Lambda} = \frac{\sqrt{2}}{3}g_{\omega N}$$

[9]. The effective mass of nucleon is

$$m_N^* = M\delta = M - g_{\sigma\Lambda}\sigma \quad (54)$$

and

$$m_\Lambda^* = M_\Lambda = M\delta_\Lambda = g_{\sigma\Lambda}\sigma - g_{\sigma*\Lambda}\sigma_* \quad (55)$$

for the  $\Lambda$  hyperon. The effective mass is presented on Fig 15. This model allows us to calculate the equation of state presented on Fig.2.




FIG. 15: The effective mass  $\delta = m_F^*/M$  for nucleon and  $\Lambda$  as function of the nucleon Fermi momentum  $x = k_F/M$ .

- 
- [1] J.Boguta, A.R.Bodmer, Nucl. Phys. **A292** (1977) 413; Serot B.D., Walecka J.D. *Recent Progress in Quantum Hadrodynamics*, Int. J. Mod. Phys. E**6**, (1997) 515-631.
  - [2] N.K. Glendenning, Z. Phys., **A327** (1987) 295; Also see in *Compact Stars* by N. K Glendenning, ( Springer-Verlag, New York, 1997).
  - [3] F. Weber *Pulsars as Astrophysical Laboratories for Nuclear and Particle Physics*, IOP Publishing, Philadelphia, (1999).
  - [4] A.W.Thomas, *Chiral symmetry and the bag model:A new starting point for nuclear physics*, Adv. Nucl. Phys. **13** (1984) 1-137; P.Guichon, Phys. Lett. **B200** (1988) 235; P. Guichon, K. Saito, E. Rodionov, A.W. Thomas, *The role of nucleon structure in finite nuclei*, <http://xxx.lanl.gov/abs/nucl-th/9509034>; H.Shen, H.Toki, *Quark mean field model for nuclear matter and finite nuclei*, <http://xxx.lanl.gov/abs/nucl-th/9911046>.
  - [5] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122** (1961) 345; 124 (1961) 246.
  - [6] M. Buballa, M. Oertel,Phys. Lett. **B457** (1999) 261; *Strange quark matter with dynamically*

- generated quark mass*, <http://xxx.lanl.gov/abs/nucl-th/9810529>.
- [7] K.Schertler, S.Leupold, J.Schaffner-Bielich, Phys.Rev. **C60** (1999) 025801; *Neutron stars and quark phases in the NLJ model*, <http://xxx.lanl.gov/abs/astro-ph/9901152>.
  - [8] P. Rehberg, S.P. Klevansky and J. Hüfner, Phys. Rev. **C53** (1996) 410.
  - [9] I. Zakout, H.J. Jaqaman, S. Pal, H.Stöcker, W. Greiner, *Hot Hypernuclear Matter in the Modified Quark Meson Coupling Model*, <http://xxx.lanl.gov/abs/nucl-th/9904084>; S. Pal, M. Hanauske, I. Zakout, H.Stöcker, W. Greiner, *Neutron star properties in the quark-meson coupling model*, <http://xxx.lanl.gov/abs/astro-ph/9905010>.
  - [10] H. Toki, U. Meyer, A. Faessler, and R. Brockmann, Phys. Rev. **C58**, 3749 (1998).
  - [11] A.R. Bodmer, Phys. Rev. **D4** (1971) 1601
  - [12] E. Witten, Phys. Rev. **D30** (1984) 272.
  - [13] E. Farhi and R.L. Jaffe, Phys. Rev. **D30** (1984) 2379.
  - [14] J. Madsen and P. Haensel (eds.), Strange Quark Matter in Physics and Astrophysics, Nucl. Phys. B (Proc. Suppl.) **24B** (1991).
  - [15] R.Manka, I.Bednarek, J.Przybyla, Phys.Rev **C62**, 15802, (2000), *The neutron star matter in a relativistic mean-field theory*, <http://xxx.lanl.gov/abs/nucl-th/0001017>.
  - [16] Y. Sugahara and H. Toki, Prog. Theo. Phys. **92** (1994) 803; H.Heiselberg, *Neutron star: recent developments*, <http://xxx.lanl.gov/abs/nucl-th/991202> (1999).
  - [17] A.L.Fetter, J.D.Walecka, *Quantum Theory of Many-Particle Systems*, McGraw-Hill, New York, (1971).
  - [18] R.Feynman, *Statistical Mechanics*, W.A.Benjamin, Inc. Reading 1972.; R.Mańka, J.Kuczyński, G.V.Vittiello, Nucl. Phys. **B276** (1986) 533; R.Mańka, Annals of Physics, G.V.Vittiello, **199** (1990) 61.
  - [19] Y. Sugahara and H. Toki, Prog. Theo. Phys. **92**, 803 (1994).
  - [20] N.K. Glendenning, J.Schaffner-Bielich, Phys. Rev. C60 (1999) 025803, <http://xxx.lanl.gov/abs/astro-ph/9810290>.
  - [21] S. Weinberg, *Gravitation and Cosmology*, John Wiley & Sons, New York, (1972).
  - [22] I. Bombaci, B.Datta, ApJ, **530** (2000) L69.
  - [23] N.K. Glendenning, Phys. Lett. **B114** (1982) 392;